Cardiology For Finals

Dr. A. Joshi

London, 3rd January 2018

SIMPLY CARDIOLOGY

Cardinal Cardiac Symptoms

- Chest pain
  - "Typical"
- Breathlessness
  - Paroxysmal Nocturnal Dyspnoea
  - Orthopnoea
- Palpitations
- Syncope/LoC
- Sudden Death (attempted)

Modifiers:
Exertional symptoms
Risk factors

SIMPLY CARDIOLOGY

Case 1

58♂

Palpitations

SIMPLY CARDIOLOGY

Case 1

History

- Sudden onset and offset
- Lasts up to a few hours
- Fast and feels irregular
- Came to A&E

- No loss of consciousness/chest pain

SIMPLY CARDIOLOGY

Case 1

PMHx

- nil

DHx

- nil

SHx

- Non-smoker
- Bottle of wine a week

FHx

- No sudden cardiac death

SIMPLY CARDIOLOGY

Case 1

Examination

- BP 120/60, HR 130, RR 16, sats 95% RA
- Irregularly irregular pulse
- No murmurs

Investigations

- Bloods + thyroid
- ECG

SIMPLY CARDIOLOGY

Case 1



SIMPLY CARDIOLOGY

Case 1



SIMPLY CARDIOLOGY

Case 1



SIMPLY CARDIOLOGY

Case 1

SIMPLY CARDIOLOGY

Case 1



SIMPLY CARDIOLOGY

Case 1



SIMPLY CARDIOLOGY

Atrial Fibrillation

Classification

Type	Definition
Paroxysmal	<24 hrs
Persistent	>24hrs Needs treatment to cardiovert
Permanent	Can’t cardiovert

SIMPLY CARDIOLOGY

Atrial fibrillation

STROKE RISK!!

- Most important thing
- More important than rhythm vs. rate management
- Really, really important.

SIMPLY CARDIOLOGY

Stroke risk

<u>CHA_2DS_2VASc score</u>

C – Congestive cardiac failure +1

H – Hypertension +1

A – Age >75yrs +2

D – Diabetes +1

S – Stroke/TIA +2

V – Vascular disease (MI, PVD) +1

A – Age >65yrs +1

Sc – Sex class, female +1

SIMPLY CARDIOLOGY

Stroke risk: CHADS-VASC

Score	Risk %/year
0	0
1	1.3
2	2.2
3	3.2
4	4.0
5	6.7
6	9.8
7	9.6
8	6.7
9	15.2

SIMPLY CARDIOLOGY

Stroke Risk - Management

- 0: Nothing
- 1: ~~Aspirin~~/Warfarin/DOAC
- >2: Warfarin/DOAC

SIMPLY CARDIOLOGY

Rate vs. Rhythm in AF

- Which is better:
  - Sinus rhythm or AF but normal heart rate?

- We don't know…
- Often patient-directed.
- Evidence and preference for sinus rhythm is increasing

SIMPLY CARDIOLOGY

"COMPROMISED" AF

When AF causes haemodynamic compromise

- Hypotensive
- Chest pain
- Signs of heart failure
- Affecting level of consciousness (GCS)
- Other markers of insufficiency (metabolic acidosis)

DC CARDIOVERSION

SIMPLY CARDIOLOGY

Rhythm vs. Rate in AF



SIMPLY CARDIOLOGY

Rate vs. Rhythm

- Why Rhythm Control?
  - Paroxysmal AF/New onset Persistent
  - Very symptomatic
  - Uncontrolled rate
  - Younger patients
  - **Mitral stenosis**
  - **Congenital Heart Disease**

SIMPLY CARDIOLOGY

Rate vs. Rhythm

- Why Rate Control?
  - Asymptomatic
  - Acceptable rate
  - Not likely to succeed
    - Long history of AF

SIMPLY CARDIOLOGY

Rhythm strategies

MUST be in AF <48hrs
OR on warfarin/NOAC for 3 weeks BEFORE/4 AFTER
OR TOE guided DCCV

- Electricity: DC Cardioversion
- Drugs
  - Flecainide (normal heart)
  - Sotalol
  - Amiodarone (last resort/short term)
- AF Ablation (not acutely)

SIMPLY CARDIOLOGY

When to do nothing?

- No symptoms
- Not “compromised” (normal obs)
- No signs of end-organ dysfunction
  - Kidneys fine, lactate fine, GCS fine
- Reversible acute illness probably the cause
  - Thyroid
  - Infection
  - Post-surgery

SIMPLY CARDIOLOGY

Rate Control Strategies

- β-blockers (bisoprolol, metoprolol)
  - Avoid in asthma
- CCA (diltiazem, verapamil)
  - Avoid in heart failure
- Digoxin
  - Careful in renal failure (K+)

- All have side-effects/toxicity states.

SIMPLY CARDIOLOGY

Case 1

- 53 yo man with palpitations
- Persistent AF (>24hrs)
- BP 120/60, HR 130, RR 16, sats 95% RA
- Irregularly irregular pulse

- Rhythm control:
  - i.v. flecainide
  - DCCV
  - Long term: β-blocker, flecainide

SIMPLY CARDIOLOGY

Case 1

- 93 yo with no symptoms
- BP 120/60, HR 70, RR 16, sats 95% RA
- Irregularly irregular pulse noted by GP

- Rate control:
  - Warfarin (unless bleeding risk)
  - β-blocker

SIMPLY CARDIOLOGY

Case 1

- 67 yo with chest pain, background of HTN.
- BP 80/30, HR 170, RR 30, sats 95% RA
- Irregularly irregular pulse

- Rhythm control:
  - Emergency DCCV
  - Warfarin

SIMPLY CARDIOLOGY

OSCE tips

- Assess the pulse → detect irregularity
  - Assess again at the carotids
  - Listen/look carefully for MITRAL signs
- "This lady has an irregularly irregular pulse, the diagnosis is most likely atrial fibrillation, but may be sinus with frequent ectopic beats."

SIMPLY CARDIOLOGY

Case 2

63 ♂

Chest pain

SIMPLY CARDIOLOGY

Chest pain

History

- S – central/sided
- O – sudden/very sudden/gradual
- C – pressure/stabbing/tearing
- R – left arm/jaw/shoulder blades
- A – nausea/vomiting/sweating/fear
- T – lasts longer than 30mins
- E – exertion/position GTN/morphine
- S – out of 10

SIMPLY CARDIOLOGY

Case 2

- History
  - Sudden onset, 2 hours ago
  - Central
  - Crushing
  - Nausea and sweatiness
  - Worse with walking
  - Got better with GTN spray
  - Feels like he's about to die

- Sometimes gets a similar pain on climbing stairs

SIMPLY CARDIOLOGY

Case 2

Medical History

- Cholesterol
- Hypertension
- Smoking (20 pack years)
- Diabetes (Type II)
- Family history

- Doesn’t like tablets.

SIMPLY CARDIOLOGY

Case 2

Drugs

- GTN spray
- Simvastatin 20mg
- Amlodipine 5mg
- Brufen 200mg prn

Social

- Father of 2
- Taxi driver
- 20 pack years smoker
- 6 pints a week (12 units)

Family

- Dad had a heart attack aged 55, now has heart failure

SIMPLY CARDIOLOGY

Case 2

Examination

- Looks ill, sweaty, clammy.
- BP 160/80, HR 100, sats 95% on 2L
- All pulses present
- No murmurs
- Clear chest

SIMPLY CARDIOLOGY

Investigations

- Chest X-ray
- ECG
- Bloods (routine)
- Troponin

SIMPLY
CARDIOLOGY



Bonus slide: Aortic Dissection

- Tearing chest pain
- Interscapular
- Missing pulses

- CT/echo
- BP <100 systolic
- Surgeons

SIMPLY CARDIOLOGY

ECGs: ACUTE MI

- Where's the clot?
- Is it a STEMI?

SIMPLY CARDIOLOGY

AMI ECG, ANATOMY AND PATHOLOGY

LCX LESIONS ±

POSTERIOR MI
STE: V7-9
STD: V1-2 (reciprocal STE)
R:S ≥ 1: V1-2
Tall T: V1-2
RCA and LCX occlusion

Seek and exclude
POSTEROLATERAL MI
STE: V7-9 and I, aVL, V5-6
STD: V1, V2
LAD and LCX occlusion

INFEROPOSTERIOR MI
STE: II, III, AVF and V7-9
STD: V1, V2 (reciprocal STE)
R:S ≥ 1: V1-2
Tall T: V1-2
RCA and LCX occlusion



RCA 'TYPE' LESIONS ±

INFERIOR MI
STE: II, III, aVF
STD: aVL (reciprocal STE)
RCA occlusion distal to RV
58% of MI

Seek and exclude
INFERIOR AND RV MI
STE: II, III, aVF and V1, V4R
RCA occlusion proximal to RV
40% of Inferior MI
Increased mortality risk

INFEROLATERAL MI
STE: II, III, AVF and I, aVL, V5, V6 ± V4R
LAD and LCX occlusion
in a L dominant system

INFEROPOSTERIOR MI
STE: II, III, AVF and V7-9
STD: V1, V2 (reciprocal STE)
R:S ≥ 1: V1-2
Tall T: V1-2
RCA and LCX occlusion

LAD LESIONS

Combinations of the following
SEPTAL MI
STE: V1-2
LAD occlusion

ANTERIOR MI
STE: V3, V4
LAD occlusion

LATERAL MI
STE: V5, V6, I, aVL
LAD occlusion

VISIBLE BODY®
© Argosy Publishing, Inc., 2007-2010
All Rights Reserved
www.visiblebody.com

Image design by Tor Ercleve
Gates, Int Med J 2005;35:263-266
3D Graphics created by Visible Body 3D
Originally posted at Lifeinthefastlane.com

PLY
IOLOGY

STEMI or NSTEMI?

Or UA?

Chest pain:
ST elevation CRITERIA
2 contiguous leads
•>1mm limb leads
•>2mm V1-V6

Chest pain:
w/other changes:
•ST depression
•T wave inversion
•Troponin!

SIMPLY CARDIOLOGY



SIMPLY CARDIOLOGY



SIMPLY CARDIOLOGY



SIMPLY CARDIOLOGY



SIMPLY CARDIOLOGY



SIMPLY
CARDIOLOGY



SIMPLY CARDIOLOGY



SIMPLY CARDIOLOGY



•Bonus slide!

<u>Pericarditis</u>

•Chest pain
•Positional
•Worse on inspiration

•Saddle shaped ST segments

•Treat with NSAIDs

•Can lead to myocarditis (trop rise)
•Heart failure

SIMPLY CARDIOLOGY

Troponin

DON'T WAIT FOR A TROPONIN IN STEMI!!



- Most sensitive at 12 hours
- But now <3 hours with high sensitivity TroponinT

Management of MI

Immediately:

- Treat pain
  - GTN
  - Morphine (co-prescribe anti-emetic)
- Give Aspirin 300mg
- $P2Y_{12}$ Inhibitor
  - Clopidogrel (300-600mg)
  - Prasugrel
  - Ticagrelor

SIMPLY CARDIOLOGY

Management of MI

STEMI

- Direct transfer for PCI
- Angiogram +/- stent
- IIb/IIIa-i (abciximab, tirofiban, eptifibatide)

NSTEMI

- "Cool off"
- Angiogram <72hrs
- Fondaparinux 2.5mg
- Risk stratify: GRACE score
- Emergency PCI if pain continues

SIMPLY CARDIOLOGY

Complications of MI

- **D**eath
- **A**rrhythmia/heart block
- **R**uptured aneurysm
- **T**hrombus (mural)
- **H**eart failure/cardiogenic shock

- **V**SD
- **A**nother MI
- **D**ressler's syndrome
- **E**moboli
- **R**egurgitant valve

OSCE Station: Median Sternotomy



- Probably a bypass
  - Look for vein graft on the legs (or it might be a LIMA only)
- Could be a valve replacement
  - Listen for metallic clicks.

SIMPLY CARDIOLOGY

Case 3

79 ♀

Breathlessness

SIMPLY CARDIOLOGY

Case 3

- At night
- Wheezy and short of breath
- "Sounded chesty"

Differential?

- Heart failure
- Pneumonia
- Asthma/COPD?
- Pulmonary embolus?!

- Orthopnoea
- Paroxysmal Nocturnal Dyspnoea (PND)
- Reducing ET (dyspnoea)

SIMPLY CARDIOLOGY

Case 3

PMHx

- MI 1999 → 2 stents
- MI 2004 → CABG
- Hypertension
- Type 2 Diabetes
- CKD 3

SIMPLY CARDIOLOGY

Case 3

DHx

- Aspirin 75mg
- Atorvastatin 40mg
- Bisoprolol 2.5mg
- Amlodipine 5mg

- New: Furosemide 40mg od from GP
  - “Swollen ankles”

SIMPLY CARDIOLOGY

Case 3

Social

- Lives alone
- Coping less well for 2/52
- No alcohol
- Ex-smoker (30py)

Family

- Parents lived to 80s
- 2 children
  - Australia
  - America

SIMPLY CARDIOLOGY

Case 3

Examination

Looks ill, cold, clammy.

BP 190/60, HR 110.

Sats 90% on RA. RR 30.

SIMPLY CARDIOLOGY

Case 3

Examination

- Raised JVP (7cm)
- Pan-systolic murmur
- Third heart sound
- Peripheral oedema to knee

Diagnosis?

- Crepitations to the midzone!

SIMPLY CARDIOLOGY

Case 3

Investigations

- Bloods
- ECG
- Chest X-ray

- Echocardiogram

Na+	128
K+	4.0
Ur	13.5
Cr	270
WCC	7
Hb	10.5
Plts	327
NT-pro-BNP	3000
CRP	<5

SIMPLY CARDIOLOGY

Case 3



SIMPLY CARDIOLOGY



IPLY
ARDIOLOGY

Heart Failure CXR ABCDE

A - Alveolar shadowing

B – Kerley-B lines

C - Cardiomegaly

D - Upper lobe Diversion

E - Pleural Effusions

SIMPLY CARDIOLOGY



IPLY
ARDIOLOGY

Heart Failure Management

Acute

- Oxygen
- Furosemide (i.v.)
  - Aiming for diuresis
- If fails → CPAP

- Off the menu!
  - GTN
  - Morphine

Long-term

- Salt restricted diet
- Cardiac rehab course

- Oral loop diuretics
- Spironolactone/Epleronone
- ACE-i/A2RB
- β-blocker
- Ivabradine

- Anti-platelets
- Defibrillator/Resynchronisation

SIMPLY CARDIOLOGY



LY
OLOGY

What causes heart failure?

- Ischaemic heart disease
- Valvular disease
- Hypertension
- Cardiomyopathy
  - Lots of rare causes

SIMPLY CARDIOLOGY

Why decompensation?

- Infarction
- Infection
- Drugs/diet
- Arrhythmia
- Thyroid disease
- Hypertension

SIMPLY
CARDIOLOGY

Heart Failure

Symptoms classification
New York Heart Association:

- NYHA I
  - No symptoms
- NYHA II
  - Mild symptoms during daily life
- NYHA III
  - Symptoms with any exertion
- NYHA IIII
  - Symptoms at rest

SIMPLY CARDIOLOGY

Heart Failure – Ejection Fraction

Mild	>45%
Moderate	35-45%
Severe	<35%

SIMPLY CARDIOLOGY

Heart Failure

Prognosis

(a)



(b)



SIMPLY CARDIOLOGY

OSCE tips

- Hard to get a decompensated patients in your exam
- Feel for a pacemaker or an ICD
- Look for RV signs without LV signs
  - JVP raised
  - Peripheral oedema
  - NO pulmonary oedema.

SIMPLY CARDIOLOGY

Case 4

88 ♂

Loss of Consciousness

SIMPLY CARDIOLOGY

Case 4

History

- Sitting at church
- Suddenly felt “wrong”
- Woke up on floor
- Immediately knew was in church
- No tongue biting/urinary incontinence/injury

SIMPLY
CARDIOLOGY

Case 4

Collateral

- Went pale
- Slumped off seat
- No shaking
- Out for “minutes”
- Normal on recovery

SIMPLY CARDIOLOGY

Case 4

PMHx
- Hypertension
- Cholesterol

DHx
- Amlodipine
- Simvastatin

SHx
- Non-smoker
- No alcohol
- Independent

FHx
- Nil

SIMPLY CARDIOLOGY

Case 4

Examination

- Looks well
- BP 190/70, HR 40, RR 16, sats 95% RA
- Ejection systolic murmur

SIMPLY CARDIOLOGY

Case 4

<u>Investigations</u>

- Bloods
- CXR
- ECG
- Echo?

SIMPLY CARDIOLOGY

ECG



SIMPLY CARDIOLOGY

ECG



SIMPLY CARDIOLOGY

ECG



SIMPLY CARDIOLOGY

Pacemakers

- 1st Degree HB
- 2nd Degree HB
  - Mobitz 1
  - Mobitz 2 (2:1, 3:1)
- Tri-fascicular block
- 3rd Degree HB (Complete)

SIMPLY CARDIOLOGY

Case 3.1

27♂

DEAD

SIMPLY CARDIOLOGY

DEAD



Torsade de Pointes

Treatment

1) Electricity
2) Magnesium (2g/10mins) (when alive)

SIMPLY CARDIOLOGY

Diagnosis?

- Drugs (cocaine, speed, MDMA)
- LQTS 1,2,3 (up to 13!)
- Brugada syndrome
- ARVC
- HCM

- All rare

SIMPLY CARDIOLOGY

Pre-excited AF/WPW



SIMPLY CARDIOLOGY

Delta-waves



SIMPLY CARDIOLOGY

Brugada



SIMPLY CARDIOLOGY

Case 5

57 ♂

Chest pain on Exertion

SIMPLY CARDIOLOGY

Case 5

History

- Worsening chest pain on exertion
- Associated with breathlessness

- Never lost consciousness

SIMPLY CARDIOLOGY

Case 5

PMHx
Nil

DHx
None

SHx

- Non-smoker
- 6 units/week
- Independent

FHx

- Nil

SIMPLY CARDIOLOGY

Case 5

- BP100/80, HR 70, sats 96%, RR 16
- Murmur heard:
  - Ejection systolic
  - Loudest over aortic region

  - Radiates to carotids.
  - Quiet second heart sound.

  - Slow rising pulse

SIMPLY CARDIOLOGY

Case 5



SIMPLY CARDIOLOGY

Case 5



SIMPLY CARDIOLOGY

Case 5



Aortic Stenosis

SIMPLY CARDIOLOGY

Aortic Valve

Tricuspid aortic valve (normal)

Bicuspid aortic valve (abnormal)



SIMPLY CARDIOLOGY

Aortic Stenosis



SIMPLY CARDIOLOGY

Aortic Stenosis

Symptoms:

- Angina
- Dyspnoea
- Syncope

Severity:

- Narrow pulse pressure
- Long murmur
- Quiet heart sound

SIMPLY
CARDIOLOGY

Aortic Stenosis

Prognosis:

- If symptomatic: poor
- 15-50% dead in a year

Treatment:

- NO GTN for angina!
- AVR (mechanical or tissue)
- TAVI

SIMPLY CARDIOLOGY

OSCE tips: Aortic Stenosis

- Unlikely to be severe in OSCE
  - So you won't get a late peaking murmur
  - Or a low volume pulse
  - But they might invent a narrow pulse pressure on the obs chart.
- More likely to get a sternotomy with a metallic second heart sound.
- Young person → bicuspid valve.

SIMPLY CARDIOLOGY

Case 6

57 ♀

Palpitations and breathlessness

SIMPLY CARDIOLOGY

Case 6

History

- 12 months of palpitations
- 3 months of exertional dyspnoea

DHx

- Amlodipine

PMHx

- Hypertension

SHx

- Independent
- Non-smoker

SIMPLY CARDIOLOGY

Case 6

Examination

140/60

HR 100 – irregularly irregular

Pan-systolic murmur, loudest at apex.

SIMPLY CARDIOLOGY

Case 6



SIMPLY CARDIOLOGY

Mitral Regurgitation

Bonus slide
Endocarditis!!



<u>Duke's criteria:</u>

MAJOR

New regurgitant valve lesion

Typical bugs on cultures

- Strep.
- Staph (post op)
- HACEK

Minor

- Vasculitic lesions
- Embolic events
- Non-typical bugs

SIMPLY CARDIOLOGY

Mitral Regurgitation

Symptoms

- Breathless on exertion
- Heart failure symptoms
- (Chest pain)
- (Palpitations)

Signs

- Pan-systolic murmur
- Atrial fibrillation
- Heart Failure

SIMPLY CARDIOLOGY

Mitral Regurgitation

Prognosis

- Acute: bad
- Chronic, symptomatic: 22% in 5 years

Treatment

Heart failure (medical) management

Anti-coagulate if in AF!

Surgery if severe MR and:

- Symptomatic w/impaired LV
- Asymptomatic w/pHTN or AF

SIMPLY CARDIOLOGY

OSCE tips

- Common in OSCEs (because it doesn't need surgery immediately)
- If you feel AF, then listen/feel for mitral disease.
  - Displaced apex
  - Pan-systolic murmur

SIMPLY CARDIOLOGY

Case 7

63 ♀

No symptoms

SIMPLY CARDIOLOGY

Case 7

- Worried about family history
  - Dad died “of MI” at 40yrs
  - Mum died of heart failure at 70yrs
  - Brother had triple bypass last year.

SIMPLY
CARDIOLOGY

Risk factor modification

- Blood pressure: 160/70mmHg
- Cholesterol: 6.3
- Diabetes: Don’t know
- Smoking: Yes
- Exercise: No

SIMPLY CARDIOLOGY

Risk Stratification: www.qrisk.org

Your results

Your risk of having a heart attack or stroke within the next 10 years is:

4.4%

In other words, in a crowd of 100 people with the same risk factors as you, 4 are likely to have a heart attack or stroke within the next 10 years.



Risk of heart attack or stroke

Your score has been calculated using the data you entered.

Your body mass index was calculated as 20.06 kg/m^2.

How does your 10-year score compare?

Your score

Your 10-year QRISK®2 score	4.4%
The score of a healthy person with the same age, sex, and ethnicity*	0.8%
Relative risk**	5.8
Your QRISK® Healthy Heart Age***	48

SIMPLY CARDIOLOGY

Risk factors

- Male sex
- Age (men ≥55 years; women ≥65 years)
- Smoking
- Dyslipidaemia
- Fasting plasma glucose 5.6–6.9 mmol/L (102–125 mg/dL)
- Abnormal Glucose TT
- Obesity
- Family history of CVD
- Asymptomatic organ damage
  - Pulse pressure (in the elderly) ≥60 mmHg
  - LVH
- Carotid wall thickening (IMT >0.9 mm) or plaque
- Carotid-femoral PWV >10 m/s
- Ankle-brachial index <0.9
- Microalbuminuria
  - Diabetes mellitus
  - Established CV or renal disease
    - Stroke
    - Ischaemic Heart Disease
- Heart failure, including heart failure with preserved EF
- Symptomatic lower extremities peripheral artery disease
- Chronic kidney disease
- Advanced retinopathy:

GIVE TABLETS

SIMPLY CARDIOLOGY

Cholesterol

Primary prevention (no heart attack, yet)

- Atorvastatin 20mg or Rosuvastatin 10mg
- Target is <40% of non-HDL cholesterol

Secondary prevention (had a heart attack)

- Atorvastatin 80mg

SIMPLY CARDIOLOGY

Summary

- MI
  - STEMI/NSTEMI?
  - Aspirin/Oxygen/Nitrates/Morphine
- Heart Failure
  - Diuretics then secondary prevention
- AF
  - Emergency?
  - Rate vs. Rhythm
- Valves
  - AS poor prognosis, early surgery
  - MR better prognosis, AF, later surgery
- Hypertension + Cholesterol
  - NICE guidelines

SIMPLY CARDIOLOGY

Cool websites

- www.lifeinthefastlane.com
- www.escardio.org (actually not that cool)
- The Simply Forum!

SIMPLY CARDIOLOGY